

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau




INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification <sup>7</sup> : <b>A61K 31/557, 9/20, 9/48</b>		A1	(11) International Publication Number: <b>WO 00/56339</b>
			(43) International Publication Date: 28 September 2000 (28.09.00)
(21) International Application Number: PCT/CA00/00301		(81) Designated States: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).	
(22) International Filing Date: 20 March 2000 (20.03.00)			
(30) Priority Data: 09/273,692 22 March 1999 (22.03.99) US			
(71) Applicant (for all designated States except US): PHARMA-SCIENCE INC. [CA/CA]; 6111 Royalmount Avenue, Suite 100, Montréal, Québec H4P 2T4 (CA).			
(72) Inventors; and (75) Inventors/Applicants (for US only): OUALI, Aomar [CA/CA]; 12 Lapointe, Boisbriand, Québec J7G 1A5 (CA). AZAD, Abul, Kalam [BD/CA]; 6495 Lavoie, Apt. #01, Montréal, Québec H3W 2K6 (CA).		Published With international search report.	
(74) Agent: SWABEY OGILVY RENAULT; Suite 1600, 1981 McGill College Avenue, Montreal, Québec H3A 2Y3 (CA).			

(54) Title: STABILIZED PHARMACEUTICAL COMPOSITION OF A NONSTEROIDAL ANTI-INFLAMMATORY AGENT AND A PROSTAGLANDIN

(57) Abstract

A pharmaceutical composition is provided for the oral administration of an NSAID and a prostaglandin. The composition is a solid dosage form wherein the NSAID is enterically coated and the prostaglandin is present along with an effective stabilizing amount of a prostaglandin stabilizing agent such as hydroxypropyl methylcellulose or polyvinylpyrrolidone. Exemplary dosage forms are bilayer tablets in which the prostaglandin is misoprostol and the NSAID is diclofenac, piroxicam, or a pharmaceutically acceptable salt thereof. Methods for using the composition to treat NSAID-responsive conditions, disorders and diseases are provided as well.



*FOR THE PURPOSES OF INFORMATION ONLY*

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Larvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
HJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LJ	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

STABILIZED PHARMACEUTICAL COMPOSITION OF A NONSTEROIDAL  
ANTI-INFLAMMATORY AGENT AND A PROSTAGLANDIN

5

TECHNICAL FIELD

This invention relates generally to pharmaceutical compositions, and more particularly relates to a pharmaceutical composition containing a combination of a nonsteroidal anti-inflammatory drug (NSAID) and a prostaglandin.

10

BACKGROUND ART

Nonsteroidal anti-inflammatory agents such as diclofenac, difenpiramide, fenbufen, flufenamic acid, ibuprofen, indomethacin, ketoprofen, meclofenamate sodium, mefenamic acid, nabumetone, naproxen, piroxicam, suprofen and tiaprofenic acid, are widely used to relieve mild to moderate pain, for fever, and to treat inflammatory conditions. Sodium diclofenac, for example, is particularly effective for relief of musculoskeletal and joint disorders such as rheumatoid arthritis, an autoimmune disease, osteoarthritis and ankylosing spondylitis; peri-acicular disorders such as bursitis and tendinitis; soft-tissue disorders such as sprains and strains, and other painful conditions such as renal colic, acute gout, dysmenorrhoea, and for relieving pain following some surgical disorders. The NSAIDs are non-habit forming drugs and thus offer a significant advantage over traditional opioid-based drugs; furthermore, as NSAIDs are by definition "nonsteroidal," the side effects commonly associated with oral administration of steroids are avoided as well. However, it is recognized that NSAIDs also exhibit some undesirable side effects, particularly at high dosages and/or with chronic oral administration. Generally, high dosages and chronic use of NSAIDs are associated with problems such as gastrointestinal and duodenal bleeding, ulceration and perforation.

In view of the advantages of NSAIDs over opioid-based drugs and steroidal agents, steps have been undertaken to minimize the drugs' adverse effects. In one approach, NSAIDs have been administered locally, such as by injection, by topical administration of, for example, an ointment or cream, by use of a transdermal patch, or by an inhalation device.

5 Although local administration is desirable, administration of an effective amount of the active agent is difficult or inconvenient. In another approach to reduce the adverse effects of NSAIDs, the agents are ingested after food or milk, or are taken in combination with antacids, histamine H<sub>2</sub>-receptor antagonists, omeprazole, or sucralfate.

10 In yet another approach to reduce the undesirable gastrointestinal effects resulting from the oral administration of NSAIDs, the agents have been co-administered with some prostaglandins, particularly "E-series" prostaglandins such as PGE<sub>1</sub>, PGE<sub>2</sub>, misoprostol, and derivatives thereof; see, e.g., U.S. Patent Nos. 3,781,429 to Partridge, 3,927,213 to Lippman, 3,928,588 to Robert, and 5,015,481 to Franz et al. Administration of a prostaglandin with an NSAID has been shown to reduce the ulcerogenicity of the NSAID. However,

15 prostaglandins are unstable compounds and degrade readily in the presence of NSAIDs, thus requiring a stabilizing agent such as hydroxypropyl methylcellulose (HPMC) or polyvinylpyrrolidone (PVP) which can, in turn, lessen the activity of an NSAID. See, for example, U.S. Patent No. 4,301,146 to Sanvordeker, which discloses prostaglandin E-type compounds stabilized with hydroxypropyl methylcellulose or polyvinylpyrrolidone before

20 being pressed into tablets, U.S. Patent No. 3,954,787 to Monkhouse, which discloses that lyophilized compositions of prostaglandin E and sodium chloride, cyclodextrin or polyvinylpyrrolidone are stable, and U.S. Patent No. 5,015,481 to Franz et al., which discusses the destabilization of prostaglandins in the presence of the NSAIDs diclofenac and piroxicam.

25 There is, accordingly, a need in the art to provide a composition for administering an NSAID wherein the undesirable gastrointestinal side effects of the drug are minimized but wherein the drug's therapeutic effectiveness is maintained. The present invention is addressed to the aforementioned need in the art and provides a stabilized pharmaceutical composition of an NSAID and a prostaglandin, i.e., a composition in which the

30 prostaglandin is stabilized and the efficacy of the NSAID is maintained.

## DISCLOSURE OF THE INVENTION

Accordingly, it is a primary object of the invention to provide a stabilized pharmaceutical composition for oral administration of an NSAID and a prostaglandin.

5 It is another object of the invention to provide such a composition wherein the NSAID is enterically coated.

It is yet another object of the invention to provide such a composition that additionally includes a prostaglandin-stabilizing agent.

10 It is still another object of the invention to provide such a composition in which the enterically coated NSAID and the prostaglandin are present in discrete regions of the composition, such as in a bilayer tablet wherein the enterically coated NSAID is present in a first layer and the prostaglandin and the prostaglandin stabilizing agent are present in a second layer.

15 Another object of the invention is to provide a method for treating a patient with an NSAID-responsive condition, disease or disorder, wherein the method comprises administering an NSAID to the patient in a stabilized pharmaceutical composition as provided herein.

20 Still another object of the invention is to provide a method for reducing the undesirable gastrointestinal side effects associated with the oral administration of an NSAID, wherein the method comprises co-administering a prostaglandin with the NSAID in a stabilized pharmaceutical composition as provided herein.

Additional objects, advantages and novel features of the invention will be set forth in part in the description which follows, and in part will become apparent to those skilled in the art upon examination of the following, or may be learned by practice of the invention.

25 In one embodiment, the first layer of the bilayer tablet comprises an enterically coated nonsteroidal anti-inflammatory agent, and the second layer comprises a prostaglandin and a stabilizing agent.

30 In another embodiment, a method of treating a patient is provided for carrying out the present therapeutic method comprising administering to the patient a pharmaceutical composition bilayer tablet as described herein.

## BRIEF DESCRIPTION OF THE DRAWINGS

Figures 1 and 2 are schematic representations of dosage forms of the invention.

5

## MODES FOR CARRYING OUT THE INVENTION

### Overview and Definitions:

Before describing the present invention in detail, it is to be understood that this  
10 invention is not limited to particular drugs or drug delivery systems, as such may vary. It is  
also to be understood that the terminology used herein is for the purpose of describing  
particular embodiments only, and is not intended to be limiting.

It must be noted that, as used in this specification and the appended claims, the  
singular forms "a," "an" and "the" include plural referents unless the context clearly dictates  
15 otherwise. Thus, for example, reference to "a pharmacologically active agent" includes a  
combination of two or more pharmacologically active agents, reference to "a stabilizer"  
includes combinations of two or more stabilizers, reference to "a prostaglandin" includes  
combinations of two or more prostaglandins, and the like.

In describing and claiming the present invention, the following terminology will be  
20 used in accordance with the definitions set out below.

The terms "active agent," "drug" and "pharmacologically active agent" are used  
interchangeably herein to refer to a chemical material or compound which, when  
administered to an organism (human or animal) induces a desired pharmacologic effect.  
Included are derivatives and analogs of those compounds or classes of compounds  
25 specifically mentioned which also induce the desired pharmacologic effect.

An "enterically coated" drug or tablet refers to a drug or tablet that is coated with a  
substance--i.e., with an "enteric coating"--that remains intact in the stomach but dissolves  
and releases the drug once the small intestine is reached.

By "pharmaceutically acceptable carrier" or "pharmaceutically acceptable vehicle"  
30 are meant materials that are suitable for oral administration and not biologically or otherwise  
undesirable, i.e., that may be administered to an individual along with an active agent

without causing any undesirable biological effects or interacting in a deleterious manner with any of the other components of the pharmaceutical composition in which it is contained.

Similarly, a "pharmaceutically acceptable" salt, ester or other derivative of an active agent as provided herein is a salt, ester or other derivative which is not biologically or otherwise undesirable.

"Stabilizing agents" as used herein refer to compounds that lower the rate at which the prostaglandins degrade, particularly in an oral pharmaceutical formulation, in the presence of an NSAID, and under environmental conditions of storage.

By "incompatible," as in two drugs that are "incompatible" with respect to each other is meant that in close physical proximity a first drug may have a deleterious effect on the physical or chemical stability of a second drug, and/or vice versa.

By the terms "effective amount" or "therapeutically effective amount" of an agent as provided herein are meant a nontoxic but sufficient amount of the agent to provide the desired therapeutic effect. As will be pointed out below, the exact amount required will vary from subject to subject, depending on the age, weight, and general condition of the subject, the severity of the condition being treated, and the like. Thus, it is not possible to specify an exact "effective amount." However, an appropriate "effective" amount in any individual case may be determined by one of ordinary skill in the art using only routine experimentation.

By a "pharmacologically acceptable" compound is meant a material which is not biologically or otherwise undesirable, i.e., the material may be administered to an individual along with the selected active agent without causing any undesirable biological effects or interacting in a deleterious manner with any of the other components of the pharmaceutical composition in which it is contained. Similarly, a "pharmacologically acceptable" salt or a "pharmacologically acceptable" ester of a compound as provided herein is a salt or ester which is not biologically or otherwise undesirable.

The invention, as noted above, is in one embodiment a stabilized pharmaceutical composition for administration of an NSAID and a prostaglandin, wherein the NSAID is enterically coated. Preferably, the composition is comprised of two discrete regions, wherein the enterically coated NSAID is present in a first region and the prostaglandin is present in a second region, along with a prostaglandin-stabilizing agent; an exemplary such

composition is a bilayer tablet as shown in Figure 1. The tablet 10 can have any geometric shape, although a generally oval shape is shown. The tablet 10 includes a first layer 11 and an adjacent second layer 12; alternatively, as shown in Figure 2, the tablet can comprise a first region 13 adjacent to a second region 14.

5       The invention is not limited with respect to the selected NSAID; the stabilized compositions of the invention can contain any NSAID, NSAID derivative, or combination of NSAIDs. Typical NSAIDs include, but are not limited to, acetylsalicylic acid, apazone, diclofenac, difenpiramide, diflunisal, etodolac, fenbufen, flufenamic acid, flurbiprofen, ibuprofen, indomethacin, ketoprofen, ketorolac, meclofenamate, mefenamic acid,  
10   nabumetone, naproxen, oxaprozin, piroxicam, sulindac, suprofen, tiaprofenic acid and tolmetin. Pharmaceutically acceptable analogs of such NSAIDs are suitable as well, particularly pharmaceutically acceptable salts. Diclofenac, piroxicam and their salts (e.g., diclofenac sodium) are particularly preferred.

      The NSAID is present in the composition in a therapeutically effective amount;  
15   preferably, the composition is in unit dosage form. The amount of NSAID administered will, of course, be dependent on the age, weight, and general condition of the subject, the severity of the condition being treated, and the judgment of the prescribing physician. Suitable therapeutic amounts will be known to those skilled in the art and/or are described in the pertinent reference texts and literature. For diclofenac sodium, for example, a  
20   therapeutic dose is typically in the range of approximately 25 mg to about 75 mg per tablet, optimally about 50 mg per tablet. The therapeutic dosing range for a tablet containing piroxicam is about 5 mg to about 50 mg per tablet, optimally about 20 mg per tablet, while the therapeutic dosing range for a tablet containing naproxen is about 250 mg to 750 mg per tablet.

25       The NSAID is enterically coated within the stabilized composition of the invention. Generally, the enteric coating comprises a polymeric material that prevents NSAID release in the low pH environment of the stomach but that ionizes at a slightly higher pH, typically a pH of 4 or 5, and thus dissolves sufficiently in the small intestines to gradually release the active agent therein. Accordingly, among the most effective enteric coating materials are  
30   polyacids having a  $pK_a$  in the range of about 3 to 5. Suitable enteric coating materials include, but are not limited to, polymerized gelatin, shellac, methacrylic acid copolymer type C NF, cellulose butyrate phthalate, cellulose hydrogen phthalate, cellulose propionate



phthalate, polyvinyl acetate phthalate (PVAP), cellulose acetate phthalate (CAP), cellulose acetate trimellitate (CAT), hydroxypropyl methylcellulose phthalate, hydroxypropyl methylcellulose acetate, dioxypopyl methylcellulose succinate, carboxymethyl ethylcellulose (CMEC), hydroxypropyl methylcellulose acetate succinate (HPMCAS), and  
5 acrylic acid polymers and copolymers, preferably formed from methyl acrylate, ethyl acrylate, methyl methacrylate and/or ethyl methacrylate with copolymers of acrylic and methacrylic acid esters (Eudragit NE, Eudragit RL, Eudragit RS) particularly preferred.

The NSAID-containing region or layer can also contain various excipients, as is well known in the pharmaceutical art, provided such excipients do not exhibit a destabilizing  
10 effect on any components in the composition. Thus, excipients such as binders, bulking agents, diluents, disintegrants, lubricants, fillers, carriers, and the like can be combined with the NSAID in the core. For solid compositions, diluents are typically necessary to increase the bulk of a tablet so that a practical size is provided for compression. Suitable diluents include dicalcium phosphate, calcium sulfate, lactose, cellulose, kaolin, mannitol, sodium  
15 chloride, dry starch and powdered sugar. Binders are used to impart cohesive qualities to a tablet formulation, and thus ensure that a tablet remains intact after compression. Suitable binder materials include, but are not limited to, starch (including corn starch and pregelatinized starch), gelatin, sugars (including sucrose, glucose, dextrose and lactose), polyethylene glycol, waxes, and natural and synthetic gums, e.g., acacia sodium alginate,  
20 polyvinylpyrrolidone, cellulosic polymers (including hydroxypropyl cellulose, hydroxypropyl methylcellulose, methyl cellulose, hydroxyethyl cellulose, and the like), and Veegum. Lubricants are used to facilitate tablet manufacture; examples of suitable lubricants include, for example, magnesium stearate, calcium stearate, and stearic acid, and are preferably present at no more than approximately 1 wt.% relative to tablet weight.  
25 Disintegrants are used to facilitate tablet disintegration or "breakup" after administration, and are generally starches, clays, celluloses, algin, gums or crosslinked polymers. If desired, the pharmaceutical composition to be administered may also contain minor amounts of nontoxic auxiliary substances such as wetting or emulsifying agents, pH buffering agents and the like, for example, sodium acetate, sorbitan monolaurate, triethanolamine sodium  
30 acetate, triethanolamine oleate, etc. If desired, flavoring, coloring and/or sweetening agents may be added as well. Other optional components for incorporation into an oral formulation herein include, but are not limited to, preservatives, suspending agents, thickening agents,

and the like. Fillers include, for example, insoluble materials such as silicon dioxide, titanium oxide, alumina, talc, kaolin, powdered cellulose, microcrystalline cellulose, and the like, as well as soluble materials such as mannitol, urea, sucrose, lactose, dextrose, sodium chloride, sorbitol, and the like.

5       The second region or layer of the composition contains a prostaglandin to reduce or eliminate the undesirable side effects of the NSAID following oral administration. Thus, preferred prostaglandins are those which are effective in this regard, i.e. are typically "anti-ulcerogenic." The prostaglandin is selected from the group consisting of naturally occurring prostaglandins, derivatives of naturally occurring prostaglandins such as hydrolyzable lower  
10   alkyl esters thereof, synthetic prostaglandins and semisynthetic prostaglandins.

      The "naturally occurring" prostaglandins useful in conjunction with the present invention are PGE<sub>0</sub>, PGE<sub>1</sub>, PGA<sub>1</sub>, PGB<sub>1</sub>, PGF<sub>1α</sub>, 19-hydroxy-PGA<sub>1</sub>, 19-hydroxy-PGB<sub>1</sub>, PGE<sub>2</sub>, PGA<sub>2</sub>, PGB<sub>2</sub>, 19-hydroxy-PGA<sub>2</sub>, 19-hydroxy-PGB<sub>2</sub>, PGE<sub>3</sub>, PGF<sub>3α</sub> and PGI<sub>2</sub>. The term "synthetic prostaglandin derivatives" is intended to encompass known or unknown  
15   compounds related to the aforementioned naturally occurring prostaglandins that are chemically synthesized using starting materials other than one of the naturally occurring prostaglandins. The term "semisynthetic prostaglandin derivatives" refers to known or unknown compounds related to the aforementioned naturally occurring prostaglandins and that are synthesized therefrom. Synthetic and semisynthetic prostaglandins include, but are  
20   not limited to, carboprost tromethamine, dinoprost tromethamine, dinoprostone, lipoprost, gemeprost, metenoprost, sulprostone and tiaprost. The preferred prostaglandin is misoprostol, present in an amount of about 50 to 500 µg per tablet, more preferably about 100 to 200 µg per tablet. Misoprostol is released immediately in the gastrointestinal tract, and produces its gastric anti-secretory effect thereby effectively reducing and/or eliminating  
25   the ulcerogenicity of the NSAID.

      The region or layer of the present pharmaceutical composition containing the prostaglandin also contains a prostaglandin stabilizing agent such as hydroxypropyl methylcellulose or polyvinylpyrrolidone, as disclosed in U.S. Patent No. 4,301,146 to Sanvorderker. Other stabilizing agents include, but are not limited to: cellulosic polymers  
30   such as hydroxypropyl cellulose, hydroxyethyl cellulose, methyl cellulose, ethyl cellulose, cellulose acetate, cellulose acetate phthalate, cellulose acetate trimellitate, hydroxypropyl methylcellulose phthalate, microcrystalline cellulose and carboxymethylcellulose sodium;

and vinyl polymers and copolymers such as polyvinyl acetate, polyvinylacetate phthalate, vinylacetate crotonic acid copolymer, and ethylene-vinyl acetate copolymers. The stabilizing agent is present in an amount effective to provide the desired stabilizing effect; generally, this means that the ratio of prostaglandin to the prostaglandin stabilizing agent is at least  
5 about 1:500 w/w, more preferably about 1:99 w/w.

The prostaglandin-containing region or layer can also contain various excipients, as discussed with respect to the NSAID-containing region or layer, i.e., excipients that do not exhibit a destabilizing effect and include, for example, binders, bulking agents, diluents, disintegrants, lubricants, fillers, carriers, and the like.

10 The active agents in the present composition, i.e., both the NSAID and the prostaglandin, may be administered in the form of a pharmacologically acceptable salt, ester, amide, prodrug or analog or as a combination thereof. Salts, esters, amides, prodrugs and analogs of the active agents may be prepared using standard procedures known to those skilled in the art of synthetic organic chemistry and described, for example, by J. March,  
15 "Advanced Organic Chemistry: Reactions, Mechanisms and Structure," 4th Ed. (New York: Wiley-Interscience, 1992). For example, basic addition salts are prepared from the neutral drug using conventional means, involving reaction of one or more of the active agent's free hydroxyl groups with a suitable base. Generally, the neutral form of the drug is dissolved in a polar organic solvent such as methanol or ethanol and the base is added thereto. The  
20 resulting salt either precipitates or may be brought out of solution by addition of a less polar solvent. Suitable bases for forming basic addition salts include, but are not limited to, inorganic bases such as sodium hydroxide, potassium hydroxide, ammonium hydroxide, calcium hydroxide, trimethylamine, or the like. Preparation of esters involves functionalization of hydroxyl groups which may be present within the molecular structure of  
25 the drug. The esters are typically acyl-substituted derivatives of free alcohol groups, i.e., moieties which are derived from carboxylic acids of the formula  $\text{RCOOH}$  where R is alkyl, and preferably is lower alkyl. Esters can be reconverted to the free acids, if desired, by using conventional hydrogenolysis or hydrolysis procedures. Preparation of amides and prodrugs can be carried out in an analogous manner. Other derivatives and analogs of the active  
30 agents may be prepared using standard techniques known to those skilled in the art of synthetic organic chemistry, or may be deduced by reference to the pertinent literature.

Before incorporation into a dosage form, the NSAID is preferably provided in a particulate form suitable for processing. Preferred types of particles are selected from the group consisting of granules, pellets, seeds and microspheres. Granules may be prepared by any art-known process. It is preferred, however, that the granules be prepared by processes  
5 such as high shear granulation, low shear granulation or fluid-bed granulation provided with top spray. Pellets or seeds can be prepared by techniques known to those skilled in the art, for example, by using a fluid-bed granulator provided with a rotor-disc. Microspheres can be prepared by any art known process with preparation by spray drying preferred.

Once in particulate form, the NSAID is enterically coated. Although any art-known  
10 process may be used, it is preferred that the enteric coating process be accomplished by utilizing either a fluid-bed coater provided with a top spray or a fluid-bed Wurster coater with a bottom spray. The resulting enterically coated particles should have a particle size in the range of about 20  $\mu\text{m}$  to 1500  $\mu\text{m}$ , preferably in the range of about 50  $\mu\text{m}$  to 300  $\mu\text{m}$ .

Before incorporation into the dosage form, the prostaglandin should be separately  
15 stabilized with the stabilizing agent. Suitable stabilization procedures are well known to those skilled in the art.

One preferred dosage form of the present invention is a bilayer tablet. Bilayer tablets as shown in Figures 1 and 2 provide several manufacturing advantages. The bilayer tablet is made in a single step compression, thereby eliminating the operations of prior methods  
20 involving first compressing one of the actives as a core tablet and subsequently coating the core, and additionally eliminating the concomitant steps of in-process and quality controls for manufacturing two different tablets. Thus, the bilayer tablet is easier and more economical to manufacture than prior compositions that separate a first drug and a second drug into physically discrete regions of a single dosage form.

25 A preferred method for forming tablets herein is by direct compression of the enterically coated NSAID and prostaglandin, optionally in combination with diluents, binders, lubricants, disintegrants, colorants or the like. As an alternative to direct compression, compressed tablets can be prepared using wet-granulation or dry-granulation processes. Tablets may also be molded rather than compressed, starting with a moist  
30 material containing a suitable water-soluble lubricant. Preferred tablets herein are manufactured using compression rather than molding, however.

In an alternative embodiment, the enterically coated NSAID and the stabilized prostaglandin are mixed into a single granulation, and the admixture is compressed into a tablet or filled into a capsule. In the admixture, there is a random possibility of the NSAID and the prostaglandin coming into contact with each other. However, the enteric coating on the NSAID granules provides a physical barrier between the NSAID and the prostaglandin, thereby minimizing direct physical contact between the two active agents. Capsule materials may be either hard or soft, and are preferably sealed, such as with gelatin bands or the like. Tablets and capsules for oral use will generally include one or more commonly used excipients as discussed earlier herein.

For administration of the dosage form, i.e., the tablet or capsule containing the enterically coated NSAID and the stabilized prostaglandin, a total weight in the range of approximately 100 mg to 1000 mg is used. The dosage form is orally administered to a patient suffering from a condition for which an NSAID would typically be indicated, including, but not limited to: alleviation of pain, e.g., arthritic pain, lumbosacral pain, musculo-skeletal pain, pain associated with a sore throat, and the like; treatment of inflammatory conditions and diseases such as osteoarthritis and rheumatoid arthritis; and treatment of psoriasis.

It is to be understood that while the invention has been described in conjunction with the preferred specific embodiments thereof, that the foregoing description as well as the examples which follow are intended to illustrate and not limit the scope of the invention. Other aspects, advantages and modifications within the scope of the invention will be apparent to those skilled in the art to which the invention pertains.

#### EXAMPLE 1

A misoprostol-HPMC complex 1% was made by mixing misoprostol with HPMC in a ratio of 1:99. Misoprostol, an oily, viscous liquid was stabilized as a solid dispersion using hydroxypropyl methycellulose as substrate and spraying misoprostol from an alcoholic solution in a fluid-bed granulator. The granules so obtained were mixed with excipients as described in Example 2.

## EXAMPLE 2

The granules of the misoprostol-HPMC complex prepared in Example 1 were mixed with the following excipients.

5

Ingredient	mg per 200 mg
Misoprostol-HPMC Complex 1%	20.0
Microcrystalline Cellulose PH 102	141.8
Crospovidone XL	8.0
Microcrystalline Cellulose PH 102	29.0
Hydrogenated Castor Oil Powder	0.8
Colloidal Silicon Dioxide	0.4

10

The blend so obtained was used in the preparation of the bilayer tablets as described in Examples 5 and 6 below.

15

## EXAMPLE 3

A blend of enterically coated granules of diclofenac was prepared as follows.

20

Ingredient	mg per 200 mg
Granulation I	
Diclofenac Sodium	50.0
Lactose Hydrous Spray Dried	15.0
Microcrystalline Cellulose PH 102	16.0
Starch (Corn) Tablet White	9.0
Povidone PVK-30	4.0
Granulation II	
Methacrylic Acid Copolymer Type C	5.4
Triacetin	0.54
Antifoam 1520-US	0.06
Microcrystalline Cellulose PH 102	98
Hydrogenated Castor Oil Powder	2

25

30

In the first step, granulation I was prepared by blending diclofenac sodium with lactose hydrous spray dried, microcrystalline cellulose PH 102, starch (corn) tablet white, and an aqueous solution of Povidone PVK-30 in a fluid-bed granulator. The granules so obtained were enteric coated in a fluid-bed granulator by the application of an enteric dispersion system containing methacrylic acid copolymer type C, NF, triacetin, and antifoam 1520-US. The enterically coated diclofenac granules were then blended with microcrystalline cellulose PH 102 and hydrogenated castor oil powder.

## EXAMPLE 4

A second composition of the diclofenac layer for the pharmaceutical delivery system of bilayer tablet was prepared consisting of enterically coated granules of diclofenac. The tablet had the following composition, and the excipients were blended according to Example 3.

Ingredients	mg per 200 mg
Granulation I	
Diclofenac Sodium	50.0
Lactose Hydrous Spray Dried	15.0
Microcrystalline Cellulose PH 102	18.0
Starch (Corn) Tablet White	9.0
Povidone PVK-30	4.0
Granulation II	
Methacrylic Acid Copolymer Type C	2.70
Triacetin	0.27
Antifoam 1520-US	0.03
Hydrogenated Castor Oil Powder	1.0

## EXAMPLE 5

A bilayer tablet was prepared containing a misoprostol solid dispersion and enterically coated granules of diclofenac. The enterically coated NSAID prepared in Example 3 was first placed in the tablet press, followed by the misoprostol blend prepared in  
5 Example 2. The tableting mechanism was then actuated to give a bilayer tablet.

## EXAMPLE 6

The misoprostol blend, prepared in Example 2, is mixed with the diclofenac blend prepared in either Example 3 or Example 4. The admixture so obtained is compressed into a  
10 tablet, or is filled into a capsule shell.



- 15 -

CLAIMS:

1. A solid pharmaceutical composition for oral administration, comprising:  
a therapeutically effective amount of a nonsteroidal anti-inflammatory  
5 drug (NSAID) coated with an enteric coating material;  
an effective anti-ulcerogenic amount of a prostaglandin; and  
an effective stabilizing amount of a prostaglandin stabilizing agent.
2. The composition of claim 1, comprising a dosage form having two discrete  
10 regions, wherein the enterically coated NSAID is present in the first of said two  
regions and the prostaglandin and prostaglandin stabilizing agent are present in  
the second of said two regions.
3. The composition of claim 2, comprising a bilayer tablet.  
15
4. The composition of claim 2, comprising a capsule.
5. The composition of claim 1, comprising an admixture of the enterically coated  
NSAID, prostaglandin and prostaglandin stabilizing agent.  
20
6. The composition of claim 5, in the form of a tablet.
7. The composition of any one of claims 1 to 6, wherein the NSAID is selected  
from the group consisting of acetylsalicylic acid, apazone, diclofenac,  
25 difenpiramide, diflunisal, etodolac, fenbufen, flufenamic acid, flurbiprofen,  
ibuprofen, indomethacin, ketoprofen, ketorolac, meclofenamate, mefenamic acid,  
nabumetone, naproxen, oxaprozin, piroxicam, sulindac, suprofen, tiaprofenic  
acid, tolmetin, pharmaceutically acceptable salts thereof, and combinations of any  
of the foregoing.

- 16 -

8. The composition of claim 7, wherein the NSAID is diclofenac or a pharmaceutically acceptable salt thereof.
- 5 9. The composition of claim 8, wherein the NSAID is diclofenac.
10. The composition of claim 8, wherein the NSAID is diclofenac sodium.
11. The composition of claim 7, wherein the NSAID is piroxicam.
- 10 12. The composition of any one of claims 1 to 11, wherein the prostaglandin is selected from the group consisting of misoprostol, PGE<sub>0</sub>, PGE<sub>1</sub>, PGA<sub>1</sub>, PGB<sub>1</sub>, PGF<sub>1</sub>, 19-hydroxy-PGA<sub>1</sub>, 19-hydroxy-PGB<sub>1</sub>, PGE<sub>2</sub>, PGA<sub>2</sub>, PGB<sub>2</sub>, 19-hydroxy-PGA<sub>2</sub>, 19-hydroxy-PGB<sub>2</sub>, PGE<sub>3</sub>, PGF<sub>3</sub>, PGI<sub>2</sub>, carboprost tromethamine, dinoprost
- 15 tromethamine, gemeprost, metenoprost, sulprostone, tiaprost and combinations thereof.
13. The composition of claim 12, wherein the prostaglandin is misoprostol.
- 20 14. The composition of any one of claims 1 to 13, wherein the prostaglandin stabilizing agent is selected from the group consisting of hydroxypropyl cellulose, hydroxyethyl cellulose, methyl cellulose, ethyl cellulose, cellulose acetate, cellulose acetate phthalate, cellulose acetate trimellitate, hydroxypropyl methylcellulose, hydroxypropyl methylcellulose phthalate, microcrystalline
- 25 cellulose, carboxymethylcellulose sodium, polyvinylpyrrolidone, polyvinyl acetate, polyvinylacetate phthalate, and vinylacetate crotonic acid copolymer, and combinations thereof.

- 17 -

15. The composition of claim 14, wherein the stabilizing agent is hydroxypropyl methylcellulose.
16. The composition of claim 14, wherein the stabilizing agent is  
5 polyvinylpyrrolidone.
17. The composition of any one of claims 1 to 16, wherein the NSAID is in particulate form.
- 10 18. The composition of claim 17, wherein the particulate form is selected from the group consisting of granules, pellets, seeds and microspheres.
19. The composition of claim 17, wherein the NSAID has a particle size in the range of about 20  $\mu\text{m}$  to 1500  $\mu\text{m}$ .  
15
20. The composition of claim 19, wherein the particle size is in the range of about 50  $\mu\text{m}$  to about 300  $\mu\text{m}$ .
21. The composition of any one of claims 1 to 20, wherein the total weight of the  
20 composition is in the range of approximately about 100 mg to about 1000 mg.
22. The composition of claim 21, wherein the amount of prostaglandin is in the range of approximately about 5  $\mu\text{g}$  to about 500  $\mu\text{g}$ .
- 25 23. A bilayer tablet for oral administration of a nonsteroidal anti-inflammatory drug (NSAID), comprising:  
(a) a first layer containing a therapeutically effective amount of an enterically coated NSAID selected from the group consisting of acetylsalicylic acid, apazone, diclofenac, difenpiramide, diflunisal, etodolac, fenbufen, flufenamic acid,

- 18 -

flurbiprofen, ibuprofen, indomethacin, ketoprofen, ketorolac, meclofenamate, mefenamic acid, nabumetone, naproxen, oxaprozin, piroxicam, sulindac, suprofen, tiaprofenic acid, tolmetin, pharmaceutically acceptable salts thereof, and combinations of any of the foregoing; and

- 5 (b) a second layer containing an effective anti-ulcerogenic amount of a prostaglandin selected from the group consisting of misoprostol, PGE<sub>0</sub>, PGE<sub>1</sub>, PGA<sub>1</sub>, PGB<sub>1</sub>, PGF<sub>1</sub>, 19-hydroxy-PGA<sub>1</sub>, 19-hydroxy-PGB<sub>1</sub>, PGE<sub>2</sub>, PGA<sub>2</sub>, PGB<sub>2</sub>, 19-hydroxy-PGA<sub>2</sub>, 19-hydroxy-PGB<sub>2</sub>, PGE<sub>3</sub>, PGF<sub>3</sub>, PGI<sub>2</sub>, carboprost tromethamine, dinoprost tromethamine, gemeprost, metenoprost, sulprostone,
- 10 tiaprost and combinations thereof, and an effective stabilizing amount of a prostaglandin stabilizing agent selected from the group consisting of hydroxypropyl cellulose, hydroxyethyl cellulose, methyl cellulose, ethyl cellulose, cellulose acetate, cellulose acetate phthalate, cellulose acetate trimellitate, hydroxypropyl methylcellulose, hydroxypropyl methylcellulose
- 15 phthalate, microcrystalline cellulose, carboxymethylcellulose sodium, polyvinylpyrrolidone, polyvinyl acetate, polyvinylacetate phthalate, and vinylacetate crotonic acid copolymer, and combinations thereof.
24. A bilayer tablet for oral administration of a nonsteroidal anti-inflammatory
- 20 drug (NSAID), comprising:
- (a) a first layer containing a therapeutically effective amount of an enterically coated NSAID selected from the group consisting of diclofenac, piroxicam, naproxen and pharmaceutically acceptable salts thereof; and
- (b) a second layer containing an effective anti-ulcerogenic amount of misoprostol
- 25 and an effective stabilizing amount of a prostaglandin stabilizing agent selected from the group consisting of hydroxypropyl methylcellulose and polyvinylpyrrolidone.

- 19 -

25. The tablet of claim 24, having a total weight in the range of approximately 100 mg to 1000 mg.

26. The tablet of claim 25, containing approximately 25 mg to 75 mg diclofenac.

5

27. The tablet of claim 25, containing approximately 5 mg to 50 mg piroxicam.

28. The tablet of claim 25, containing approximately 250 mg to 750 mg naproxen.

10

29. The tablet of claim 25, containing approximately 100 µg to 200 µg misoprostol.

30. A tablet comprising an admixture of (a) a therapeutically effective amount of an enterically coated NSAID selected from the group consisting of diclofenac, piroxicam and pharmaceutically acceptable salts thereof; (b) an effective anti-ulcerogenic amount of misoprostol; and (c) an effective stabilizing amount of a prostaglandin stabilizing agent selected from the group consisting of hydroxypropyl methyl cellulose and polyvinylpyrrolidone.

15

20

31. A method for treating a patient suffering from a condition, disease or disorder that is responsive to an NSAID, comprising orally administering to the patient the pharmaceutical composition of any one of claims 1 to 22 or the tablet of any one of claims 23 to 30.

25

32. The method of claim 31, wherein the composition or tablet is administered twice daily.




Figure 1




Figure 2

# INTERNATIONAL SEARCH REPORT

Inter national Application No

PCT/CA 00/00301

## A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 A61K31/557 A61K9/20 A61K9/48

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

WPI Data, PAJ, EPO-Internal

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 91 16895 A (G. D. SEARLE) 14 November 1991 (1991-11-14)  claims 1-3,7-11 examples 2-9	1,2, 4-16,21, 22,30-32
P,X	WO 00 01368 A (NORTON HEALTHCARE) 13 January 2000 (2000-01-13) claims 1,3,5,16,19,24,26 page 2, paragraph 3 page 3, paragraph 3 page 4, paragraph 1 page 5; example 3  -/-	1-3,5-32

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

\* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

29 June 2000

Date of mailing of the international search report

06/07/2000

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
Fax: (+31-70) 340-3016

Authorized officer

Ventura Amat, A

# INTERNATIONAL SEARCH REPORT

Inter national Application No  
PCT/CA 00/00301

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P, X	WO 99 65496 A (SHERMAN, BERNARD CHARLES) 23 December 1999 (1999-12-23)  claims 1-3 page 5; examples 1-4 ----	1,2, 5-15,21, 22,31,32
E	WO 00 15200 A (NORTON HEALTHCARE) 23 March 2000 (2000-03-23)  claims 1-3,7,8,16 page 5; examples 1,2 ----	1,2, 4-15, 17-22
A	WO 99 12524 A (NYCOMED DANMARK) 18 March 1999 (1999-03-18) the whole document -----	1-32



# INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/CA 00/00301

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 9116895	A	14-11-1991	AT 121625 T	15-05-1995
			AU 7876491 A	27-11-1991
			CA 2082944 A,C	04-11-1991
			DE 69109286 D	01-06-1995
			DE 69109286 T	28-09-1995
			DK 527887 T	03-07-1995
			EP 0527887 A	24-02-1993
			ES 2071312 T	16-06-1995
			GR 3015939 T	31-07-1995
			IE 68596 B	26-06-1996
			IL 98033 A	31-07-1995
			NZ 238024 A	26-08-1993
			PT 97562 A,B	28-02-1992
			US 5601843 A	11-02-1997
			US 5698225 A	16-12-1997
			ZA 9103353 A	28-04-1993
WO 0001368	A	13-01-2000	AU 4519399 A	24-01-2000
WO 9965496	A	23-12-1999	AU 4127599 A	05-01-2000
WO 0015200	A	23-03-2000	AU 5750799 A	03-04-2000
WO 9912524	A	18-03-1999	AU 9062998 A	29-03-1999
			NO 20001290 A	28-04-2000